



## Research Article

# The Historical/Evolutionary Cause and Possible Treatment of Pandemic COVID-19 (SARS-CoV-2, 2019-CORONA VIRUS): World-War III: The Blackout of the Modern World by Neglected Small Infectious Agent

Sorush Niknamian<sup>1\*</sup> and Somayeh Zaminpira<sup>2</sup>

<sup>1</sup>Military Medicine Department, American Military University and Liberty University, USA

<sup>2</sup>Phycology and Psychiatry Department, American Beirut University, Beirut, Lebanon

## Abstract

**Background:** A virus is a small infectious agent that replicates only inside the living cells of an organism. Viruses can infect all types of life forms, from animals and plants to microorganisms, including bacteria and *Archaea*. In evolution, viruses are an important means of horizontal gene transfer, which increases genetic diversity in a way analogous to sexual reproduction. Influenza (Including COVID-19), is an infectious disease caused by an influenza virus. Some viruses especially smallpox, throughout history, has killed between 300-500 million people in its 12,000-year existence. As modern humans increased in numbers, new infectious diseases emerged, including SARS-CoV-2. We have two groups of virus, RNA and DNA viruses. The most brutal viruses are RNA ones like COVID-19 (Sars-CoV-2). Introduction: Corona viruses are a group of viruses that cause diseases in mammals and birds. In humans, corona viruses cause respiratory tract infections that are typically mild, such as some cases of the common cold (among other possible causes, predom-

inantly rhinoviruses), though rarer forms can be lethal, such as SARS, MERS and COVID-19. Symptoms vary in other species: in chickens, they cause an upper respiratory tract disease, while in cows and pigs they cause diarrhea. Corona viruses constitute the subfamily Orthocoronavirinae. The genome size, corona viruses ranges from approximately 27 to 34 kilo bases, the largest among known RNA viruses.

**Discussion and Results:** We have researched from the first virus in the planet to the last mutated version which is SARS-COV-2. We have collected many informative data in tables and figures to reach the main cause of 2019-Coronavirus and calculated the probability and estimated deaths in the current time. We have discussed about the possible treatment and prevention of the virus and did algebraic calculations on the epidemiology, the size and even the future of this pandemic. The only era which any virus had not been epidemic, were through world war 2, were the German scientists had found the way to fight any viral infections which is very important and can help scientists to reach the main treatment of the new 2019-Coronavirus. We have sorted the deadly and non-deadly corona viruses and explained how this epidemic had begun through Evolutionary Medicine (EM). The result of the article is that 16% of the whole population in the world has been contaminated which is 1248000000 of 7.8 billion people world-wide. SARS-CoV-2 is an RNA virus. Its nucleic acid is single-stranded RNA (ssRNA). The polarity of this virus is positive-sense ((+) ssRNA). Positive-sense viral RNA is similar to mRNA and thus can be immediately translated by the host cell. Recombination in RNA viruses appears to be an adaptation for coping with genome damage. Recombination can occur infrequently between animal viruses of the same species but of divergent lineages. The resulting recombinant viruses may sometimes cause an outbreak of infection in humans. RNA viruses have very high mutation rates this is one reason why it is difficult to make effective vaccines to prevent diseases caused by RNA viruses. The resulting recombinant viruses causes an outbreak of infection in humans.

**Conclusion:** In conclusion, the mutation of the SARS-CoV and influenza viruses through Drift and Reassortment is the main cause of SARS-CoV-2 through natural selection, Lamarckian Evolution and co evolution which caused this RNA virus so powerful, unpredictable and different in the genome size and nations worldwide. The first Pandemic of Influenza was first detected in 1732 and this virus evolved through natural selection till 2019 which caused the world-wide pandemic of SARS-CoV-2. Based on many studies, inhalation of Ozone plus Sulfur Dioxide, increasing the amounts of L-Glutathione (Which is low in children and older adults and this is the main reason why older adults and children die from this disease.) plus Viral Phage Therapy (VPT) which we discussed fully in this article can be the possible prime treatment of SARS-CoV-2 infection. The seasonal temperature cannot be useful in controlling/reducing the pandemic of this virus since the natural selection, Lamarckian Evolution and high mutation of the virus helps its survival. No antiviral drugs will be useful against SARS-CoV-2 because of high rate of mutation and primarily adaptation of the virus to the drugs and even the environmental Temperature.

**Keywords:** COVID-19 (SARS-CoV-2); Drift; Fibonacci Series; Influenza; L-Glutathione; Lamarckian Evolution; Natural Selection; Ozone; Positive-Sense ((+) ssRNA), Reassortment; Pandemic; Sulfur Dioxide; Viral Phage Therapy

\*Corresponding author: Sorush Niknamian, Military Medicine Department, Liberty University, USA, Tel: +1 5102813388; E-mail: so.niknamian@gmail.com, niknamian@mail.co.uk

**Citation:** Niknamian S, Zaminpira S (2020) The Historical/Evolutionary Cause and Possible Treatment of Pandemic COVID-19 (SARS-CoV-2, 2019-CORONAVIRUS): World-War III: The Blackout of the Modern World by Neglected Small Infectious Agent. J Altern Complement Integr Med 6: 099.

**Received:** April 03, 2020; **Accepted:** April 08, 2020; **Published:** April 15, 2020

**Copyright:** © 2020 Niknamian S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction


Brief history of viruses

Prehistory

The world, new infectious diseases emerged, including those caused by viruses [1]. Earlier, humans lived in small, isolated communities and most epidemic diseases did not exist [2,3]. Smallpox, which is the most lethal and devastating viral infection in history, first emerged among agricultural communities in India about 11,000 years ago [4]. The virus, which only infected humans, probably descended from the poxviruses of rodents [5]. Humans probably came into contact with these rodents, and some people became infected by the viruses they carried. When viruses cross this so-called species barrier, their effects can be severe and humans may have had little natural resistance [6]. Contemporary humans lived in small communities, and those who succumbed to infection either died or developed immunity. This acquired immunity is only passed down to offspring temporarily, by antibodies in breast milk and other antibodies that cross the placenta from the mother's blood to the unborn child's. Therefore, sporadic outbreaks probably occurred in each generation. In about 9000 BC, when many people began to settle on the fertile flood plains of the River Nile, the population became dense enough for the virus to maintain a constant presence because of the high concentration of susceptible people [7]. Other epidemics of viral diseases that depend on large concentrations of people, such as mumps, rubella and polio, also first occurred at this time [8]. The Neolithic age, which began in the Middle East in about 9500 BC, was a time when humans became farmers [9]. This agricultural revolution embraced the development of monoculture and presented an opportunity for the rapid spread of several species of plant viruses [10]. The divergence and spread of sobemoviruses southern bean mosaic virus date from this time [11]. The spread of the potyviruses of potatoes, and other fruits and vegetables, began about 6,600 years ago [10]. About 10,000 years ago the humans who inhabited the lands around the Mediterranean basin began to domesticate wild animals. Pigs, cattle, goats, sheep, horses, camels, cats and dogs were all kept and bred in captivity [12]. These animals would have brought their viruses with them [13]. The transmission of viruses from animals to humans can occur, but such zoonotic infections are rare and subsequent human-to-human transmission of animal viruses is even rarer, although there are notable exceptions such as influenza. Most viruses are species-specific and would have posed no threat to humans [14]. The rare epidemics of viral diseases originating in animals would have been short-lived because the viruses were not fully adapted to humans and the human populations were too small to maintain the chains of infection [15,16]. Other, more ancient, viruses have been less of a threat. Herpes viruses first infected the ancestors of modern humans over 80 million years ago [17]. Humans have developed a tolerance to these viruses, and most are infected with at least one species [18]. Records of these milder virus infections are rare, but it is likely that early hominids suffered from colds, influenza and diarrhoea caused by viruses just as humans do today. More recently evolved viruses cause epidemics and pandemics [17]. The influenza virus is one that seems to have crossed the species barrier from ducks and water fowl to pigs and from there to humans. It is possible that a fatal plague in the Middle East at the time of the late 18th Dynasty was associated with this transmission at Amarna [19,20].

Influenza

Influenza, commonly known as the flu, is an infectious disease caused by an influenza virus [21]. Symptoms can be mild to severe [22]. The most common symptoms include: high fever, runny nose, sore throat, muscle and joint pain, headache, coughing and feeling tired just like Sars-CoV-2 [21]. These symptoms typically begin two days after exposure to the virus and most last less than a week [21]. The cough, however, may last for more than two weeks [21]. In children, there may be diarrhea and vomiting, but these are not common in adults [23]. Diarrhea and vomiting occur more commonly in gastroenteritis, which is an unrelated disease and sometimes inaccurately referred to as stomach flu or the 24-hour flu [23]. Complications of influenza may include viral pneumonia, secondary bacterial pneumonia, sinus infections, and worsening of previous health problems such as asthma or heart failure [22,24]. Three of the four types of influenza viruses affect humans: Type A, Type B and Type C [24,25]. Type D has not been known to infect humans, but is believed to have the potential to do so [25,26]. Usually, the virus is spread through the air from coughs or sneezes [21]. This is believed to occur mostly over relatively short distances [27]. It can also be spread by touching surfaces contaminated by the virus and then touching the eyes, nose, or mouth [22,27,28]. A person may be infectious to others both before and during the time they are showing symptoms [22]. The infection may be confirmed by testing the throat, sputum, or nose for the virus [24]. A number of rapid tests are available; however, people may still have the infection even if the results are negative [24]. A type of polymerase chain reaction that detects the virus's RNA is more accurate [24] (Figure 1).






**Figure 1:** Influenza virus, magnified approximately 100,000 times, Specialty: Infectious disease, Symptoms: Fever, runny nose, sore throat, muscle and joint pain, headache, coughing, feeling tired, Usual onset: One to four days after exposure, Duration: 1 week, Causes: Influenza viruses.

**Prevention:** Hand washing, influenza vaccine, surgical masks, Frequency: 3-5 million severe cases per year. Deaths: Up to 650,000 respiratory deaths per year [1-3] (Table 1 & Figures 2-4 ).

Symptom:	Sensitivity	Specificity
Fever	68–86%	25–73%
Cough	84–98%	7–29%
Nasal congestion	68–91%	19–41%

**Table 1:** All three findings, especially fever, were less sensitive in people over 60 years of age.

Most sensitive symptoms for diagnosing influenza [1-3] .



### List of Epidemics of Plagues (Tables 2-7)

Corona viruses are a group of viruses that cause diseases in mammals and birds. In humans, corona viruses cause respiratory tract

infections that are typically mild, such as some cases of the common cold (among other possible causes, predominantly rhinoviruses), though rarer forms can be lethal, such as SARS, MERS, and COVID-19 (Tables 2-7). Symptoms vary in other species: in chickens, they cause an upper respiratory tract disease, while in cows and pigs they cause diarrhea. Corona viruses constitute the subfamily Orthocoronavirinae, in the family Coronaviridae, order Nidovirales and realm Ribavirin [39,186]. They are enveloped viruses with a positive-sense single-stranded RNA genome and a nucleocapsid of helical symmetry. The genome size of corona viruses ranges from approximately 27 to 34 kilo bases, the largest among known RNA viruses [34].

### Evolution

The most recent common ancestor (MRCA) of all corona viruses has been placed at around 8000 BCE [187]. The MRCAs of the Alpha-corona virus line has been placed at about 2400 BCE, the Beta-corona virus line at 3300 BCE, the Gamma-corona virus line at 2800 BCE, and the Delta-corona virus line at about 3000 BCE. It appears that bats and birds, as warm-blooded flying vertebrates, are ideal hosts for the corona virus gene source (with bats for Alpha-corona virus and Beta corona virus and birds for Gamma-corona virus and Delta-corona virus) to fuel corona virus evolution and dissemination [188]. Bovine corona virus and canine respiratory corona viruses diverged from a common ancestor in 1951 [189]. Bovine corona virus and human corona virus OC43 diverged around the 1890s. Bovine corona virus diverged from the equine corona virus species at the end of the 18th century [190]. The MRCA of human corona virus OC43 has been dated to the 1950s [191]. MERS-CoV, although related to several bat corona virus species, appears to have diverged from these several centuries ago [192]. The human corona virus NL63 and a bat corona virus shared an MRCA 563–822 years ago [193]. The most closely related bat corona virus and SARS-CoV diverged in 1986 [194]. A path of evolution of the SARS virus and keen relationship with bats have been proposed. The corona viruses have been coevolved with bats for a long time and the ancestors of SARS-CoV first infected the species of the genus Hipposideridae, subsequently spread to species of the Rhinolophidae and then to civets, and finally to humans [195,196]. Alpaca corona virus and human corona virus 229E diverged before 1960 [197].

### Discussion and Results

By mentioning all the facts, evolution and all the similarities from the viruses, we observe that all kinds of Flu including COVID-19 which have been caused many deaths through epidemic and pandemic of these giant but very small objects, all the viruses from the first one before the rise of the human in evolution, had been existed. Through my deep research, the first virus was or blue-green algae, pre-date viruses and are one of the oldest forms of life on Earth (Figure 5).

The researchers developed algorithms to compare the protein shapes of 3,460 viruses and 1,620 cells. They found that 442 protein folds were shared between cells and viruses, but 66 folds were unique to viruses. To make sense of the data, the team arranged the protein folds into a tree that grew a new 'branch' every time a new type of protein fold evolved. Wherever possible, the team used fossil evidence to put an approximate date on the budding of specific branches. For example, one particular protein fold was first seen in cyanobacteria (blue-green algae) and later appeared in all its descendants.

Death toll (estimate)	Location	Date	Event	Disease
75,000–100,000	Greece	429–426 BC	Plague of Athens	Unknown, possibly typhus, typhoid fever or viral hemorrhagic fever
5–10 million	Roman Empire	165–180 (possibly up to 190)	Antonine Plague	Unknown, possibly smallpox
	Europe	250–266	Plague of Cyprian	Unknown, possibly smallpox
25–50 million; 40% of population	Europe, Egypt and West Asia	541–542	Plague of Justinian	Plague
	Rome	590	Roman Plague of 590	Plague
> 100,000	Ctesiphon, Persia	627–628	Plague of Sheroe	Plague
	British Isles	664–689	Plague of 664	Plague
	Japan	735–737	735–737 Japanese small-pox epidemic	Smallpox
	Byzantine Empire, West Asia, Africa	746–747	Plague of 746–747	Plague
20–200 million; 20–60% of European population	Europe, Asia and North Africa	1331–1353	Black Death	PlagueY. pestis
> 10,000	England and later continental Europe	1485–1551	Sweating sickness (multiple outbreaks)	Unknown, possibly an unknown species of hantavirus

**Table 2:** 15th century and earlier.

Death toll (estimate)	Location	Date	Event	Disease	Ref.
5–15 million (80% of population)	Mexico	1545–1548	Cocoliztli Epidemic of 1545–1548	Possibly <i>Salmonella enterica</i>	[29,30,31]
2–2.5 million (50% of population)	Mexico	1576–1580	Cocoliztli epidemic of 1576	Possibly <i>Salmonella enterica</i>	[29,30,31,32]
280,000	Italy	1629–1631	Italian plague of 1629–1631	Plague	[33]
100,000 [34]	England	1665–1666	Great Plague of London	Plague	[35]
76,000	Austria	1679	Great Plague of Vienna	Plague	
40,000	France	1668		Plague	[36]
30–90% of population	Southern New England, especially the Wampanoag people	1616–1619	1616 New England epidemic	Unknown cause. Latest research suggests epidemic(s) of leptospirosis with Weil syndrome. Classic explanations include yellow fever, bubonic plague, influenza, smallpox, chickenpox, typhus, and syndemic infection of hepatitis B and hepatitis D.	[37,38]
24,148[39]	Netherlands	1663–1664		Plague	
> 20,100 in London	London	1563–1564	1563 London plague	Plague	
> 19,900 in London and outer parishes	London	1592–1593	1592–93 London plague	Plague	
	Seneca nation	1592–1596		Measles	[40]
	Spain	1596–1602		Plague	[41]
	South America	1600–1650		Malaria	[42]
	England	1603		Plague	[43]
	Egypt	1609		Plague	
	Wyandot people	1634		Smallpox	[44]
	Thirteen Colonies	1633	Massachusetts smallpox epidemic	Smallpox	
	England	1636		Plague	[45]
	China	1641–1644		Plague	[46]
	Spain	1647–1652	Great Plague of Seville	Plague	
	Central America	1648		Yellow fever	[47]
	Italy	1656	Naples Plague	Plague	[48]
	Thirteen Colonies	1657		Measles	[49]
	Spain	1676–1685		Plague	[35]
	Thirteen Colonies	1687		Measles	[36]
	Thirteen Colonies	1690		Yellow fever	

**Table 3:** 16–17th centuries.



	Canada, New France	1702–1703		Smallpox	[50]
> 18,000 (36% of population)	Iceland	1707–1709	Great Smallpox Epidemic	Smallpox	
	Denmark, Sweden	1710–1712	Great Northern War plague outbreak	Plague	
	Thirteen Colonies	1713–1715		Measles	[51]
	Canada, New France	1714–1715		Measles	[52]
	France	1720–1722	Great Plague of Marseille	Plague	[53]
	Thirteen Colonies	1721–1722		Smallpox	[54]
	Thirteen Colonies	1729		Measles	[55]
	Spain	1730		Yellow fever	
	Thirteen Colonies	1732–1733		Influenza	[56]
	Canada, New France	1733		Smallpox	[57]
> 50,000	Balkans	1738	Great Plague of 1738	Plague	
	Thirteen Colonies	1738		Smallpox	[58]
	Thirteen Colonies	1739–1740		Measles	
	Italy	1743		Plague	[59]
	Thirteen Colonies	1747		Measles	
	North America	1755–1756		Smallpox	
	North America	1759		Measles	[60]
	North America, West Indies	1761		Influenza	
	North America, present-day Pittsburgh area.	1763		Smallpox	[61]
> 50,000	Russia	1770–1772	Russian plague of 1770–1772	Plague	
	Pacific Northwest natives	1770s		Smallpox	[62]
	North America	1772		Measles	
> 2,000,000	Persia	1772	1772 Persia plague	Plague	[8]
	England	1775–1776		Influenza	[63]
	Spain	1778		Dengue fever	[64]
	Plains Indians	1780–1782	North American smallpox epidemic	Smallpox	[65]
	Pueblo Indians	1788		Smallpox	[66]
	United States	1788		Measles	
	New South Wales, Australia	1789–1790		Smallpox	[67]
	United States	1793		Influenza and Epidemic Typhus	
	United States	1793–1798	Yellow Fever Epidemic of 1793, resurgences	Yellow fever	[68]

**Table 4:** 18th century.

Death toll (estimate)	Location	Date	Event	Disease	Ref.
1,000,000	Russia	1852–1860	Third cholera pandemic	Cholera	[69]
1,000,000	Worldwide	1889–1890	1889–1890 flu pandemic	Influenza	[64]
> 100,000	Asia, Europe	1816–1826	First cholera pandemic	Cholera	[70]
> 100,000	Asia, Europe, North America	1829–1851	Second cholera pandemic	Cholera	[70]
40,000	Fiji	1875	1875 Fiji Measles outbreak	Measles	[62]
> 20,000	Canada	1847–1848	Typhus epidemic of 1847	Epidemic typhus	[50]
> 9,000	India, Germany	1881–1896	Fifth cholera pandemic	Cholera	[70]
4,737	Copenhagen, Denmark	1853	Cholera epidemic of Copenhagen 1853	Cholera	[54]
3,164	Montreal	1885		Smallpox	timeline
> 3,000	Central Coast, British Columbia	1862–1863		Smallpox	[59]
616	England	1854	Broad Street cholera outbreak	Cholera	[55]
	Spain	1800–1803		Yellow fever	[67]
	Ottoman Empire, Egypt	1801		Bubonic plague	[68]

	United States	1803		Yellow fever	
	Egypt	1812		Plague	
	Ottoman Empire	1812		Plague	[69]
	Malta	1813		Plague	
	Romania	1813	Caragea's plague	Plague	
	Ireland	1816–1819		Typhus	
	United States	1820–1823		Yellow fever	
	Spain	1821		Yellow fever	[71]
	New South Wales, Australia	1828		Smallpox	[72]
	Netherlands	1829	Groningen epidemic	Malaria	
	South Australia	1829		Smallpox	[73]
	Iran	1829–1835		Bubonic plague	[74]
	Egypt	1831		Cholera	[75,76]
	Plains Indians	1831–1834		Smallpox	
	England, France	1832		Cholera	
	North America	1832		Cholera	[77]
	United States	1833		Cholera	
	United States	1834		Cholera	
	Egypt	1834–1836		Bubonic plague	[77,78]
	United States	1837		Typhus	
	Great Plains	1837–1838	1837–38 smallpox epidemic	Smallpox	[79]
	Dalmatia	1840		Plague	
	South Africa	1840		Smallpox	
	United States	1841		Yellow fever	
	United States	1847		Yellow fever	
	Worldwide	1847–1848		Influenza	[80]
	Egypt	1848		Cholera	[77,78]
	North America	1848–1849		Cholera	
	United States	1850		Yellow fever	
	North America	1850–1851		Influenza	
	United States	1851		Cholera	[81]
	United States	1852		Yellow fever	
	Ottoman Empire	1853		Plague	[82]
	United States	1855		Yellow fever	
	Worldwide	1855–1960	Third plague pandemic	Bubonic plague	[83]
	Portugal	1857		Yellow fever	
	Victoria, Australia	1857		Smallpox	[84]
	Europe, North America, South America	1857–1859		Influenza	[85]
	Middle East	1863–1879	Fourth cholera pandemic	Cholera	[86]
	Egypt	1865		Cholera	[77,78]
	Russia, Germany	1866–1867		Cholera	
	Australia	1867		Measles	
	Iraq	1867		Plague	[87]
	Argentina	1852–1871		Yellow fever	[88]
	Germany	1870–1871		Smallpox	
	Russian Empire	1877		Plague	[89]
	Egypt	1881		Cholera	[77,78]
	West Africa	1900		Yellow fever	

**Table 5:** 19th century.

Death toll (estimate)	Location	Date	Event	Disease	Ref.
	Congo Basin	1896–1906		Trypanosomiasis	[90]
> 800,000	Europe, Asia, Africa	1899–1923	Sixth cholera pandemic	Cholera	[66]
113	San Francisco	1900–1904		Bubonic plague	[91]
	Uganda	1900–1920		Trypanosomiasis	[92]
	Egypt	1902		Cholera	[77,78]
22	India	1903		Bubonic Plague	[93]
4	Fremantle	1903		Bubonic plague	[94]
40,000	China	1910–1912	1910 China plague	Bubonic plague	[95]
1.5 million	worldwide	1915–1926	1915 Encephalitis lethargica pandemic	Encephalitis lethargica	[96]
up to 100,000,000	worldwide	1918–1920	Spanish flu	Influenza Spanish Flu Virus	[97]
	Russia	1918–1922		Typhus	
30	Los Angeles	1924	1924 Los Angeles pneumonic plague outbreak	Pneumonic plague	
43	Croydon, UK	1937	Croydon epidemic of typhoid fever	Typhoid fever	[98]
	Egypt	1942–1944		Malaria	[77,78]
	China	1946		Bubonic plague	
	Egypt	1946		Relapsing fever	[77,78]
	Egypt	1947		Cholera	[77,78]
2,000,000	worldwide	1957–1958	Asian flu	Influenza	[99]
	worldwide	1961–1975	Seventh cholera pandemic	Cholera	[91]
4	Sweden	1963		Smallpox	[100,101]
1,000,000	worldwide	1968–1969	Hong Kong flu	Influenza	[99]
5	Netherlands	1971		Poliomyelitis	[102]
35	Yugoslavia	1972	1972 outbreak of smallpox in Yugoslavia	Smallpox	
	United States	1972–1973	London flu	Influenza	[103]
15,000	India	1974	1974 smallpox epidemic of India	Smallpox	[104]
> 32,000,000	worldwide (commenced in Congo Basin)	1960–present	HIV/AIDS pandemic	HIV/AIDS	[105]
	South America	1990s		Cholera	
52	India	1994	1994 plague epidemic in Surat	Plague	[106]
231	worldwide	1996–2001		vCJD	
	West Africa	1996		Meningitis	
105	Malaysia	1998–1999	1998–99 Malaysia Nipah virus outbreak	Nipah virus infection	[107]
	Central America	2000		Dengue fever	[108]

**Table 6:** 20th century.

Death toll (estimate)	Location	Date	Event	Disease	Ref.
> 400	Nigeria	2001		Cholera	[109]
	South Africa	2001		Cholera	[110]
349	China	2002–2004	SARS outbreak	SARS coronavirus	[111]
299	Hong Kong	2002–2004	SARS outbreak	SARS coronavirus	[112]
37	Taiwan	2002–2004	SARS outbreak	SARS coronavirus	[113]
44	Canada	2002–2004	SARS outbreak	SARS coronavirus	[114]
33	Singapore	2002–2004	SARS outbreak	SARS coronavirus	[115]

	Algeria	2003		Plague	[116]
	Afghanistan	2004		Leishmaniasis	[117]
	Bangladesh	2004		Cholera	[118]
	Indonesia	2004		Dengue fever	[119]
	Senegal	2004		Cholera	[120]
7	Sudan	2004		Ebola	[121]
	Mali	2005		Yellow fever	[122]
27	Singapore	2005	2005 dengue outbreak in Singapore	Dengue fever	[123]
	Luanda, Angola	2006		Cholera	[124]
61	Ituri Province, Democratic Republic of the Congo	2006		Plague	[125][126]
17	India	2006		Malaria	[127]
> 50	India	2006	2006 dengue outbreak in India	Dengue fever	[127]
	India	2006	Chikungunya outbreaks	Chikungunya virus	[128]
> 50	Pakistan	2006	2006 dengue outbreak in Pakistan	Dengue fever	[129]
	Philippines	2006		Dengue fever	[130]
187	Democratic Republic of the Congo	2007	Mwekaebola epidemic	Ebola	[131]
	Ethiopia	2007		Cholera	[132]
49	India	2008		Cholera	[133]
10	Iraq	2007	2007 Iraq cholera outbreak	Cholera	[134]
	Nigeria	2007		Poliomyelitis	[135]
	Puerto Rico; Dominican Republic; Mexico	2007		Dengue fever	[136]
	Somalia	2007		Cholera	[137]
37	Uganda	2007		Ebola	[138]
	Vietnam	2007		Cholera	[139]
	Brazil	2008		Dengue fever	[140]
	Cambodia	2008		Dengue fever	[141]
	Chad	2008		Cholera	[142]
	China	2008–2017		Hand, foot and mouth disease	[143]
	Madagascar	2008		Bubonic plague	[144]
	Philippines	2008		Dengue fever	[145]
	Vietnam	2008		Cholera	[146]
4,293	Zimbabwe	2008–2009	2008–2009 Zimbabwean cholera outbreak	Cholera	[147]
18	Bolivia	2009	2009 Bolivian dengue fever epidemic	Dengue fever	[148]
49	India	2009	2009 Gujarat hepatitis outbreak	Hepatitis B	[149]
	Queensland, Australia	2009		Dengue fever	[150]
	Worldwide	2009	Mumps outbreaks in the 2000s	Mumps	
931	West Africa	2009–2010	2009–2010 West African meningitis outbreak	Meningitis	[151]
203,000	Worldwide	2009	2009 flu pandemic	Influenza	[152,153]
9,985 (May 2017)	Hispaniola	2010–present	Haiti cholera outbreak	Cholera	[154,155]
> 4,500 (February 2014)	Democratic Republic of the Congo	2011–present		Measles	[156,157]
170	Vietnam	2011–present		Hand, foot and mouth disease	[158,159]
> 350	Pakistan	2011	2011 dengue outbreak in Pakistan	Dengue fever	[160]
171 (as of 10 January 2013 <sup>[update]</sup> )	Darfur Sudan	2012	2012 yellow fever outbreak in Darfur, Sudan	Yellow fever	[161]
862 (as of 13 January 2020 <sup>[update]</sup> )	Worldwide	2012–present	2012 Middle East respiratory syndrome coronavirus outbreak	Middle East respiratory syndrome	[162,163,164]
142	Vietnam	2013–2014		Measles	[165]
>> 11,300	Worldwide, primarily concentrated in West Africa	2013–2016	Ebola virus epidemic in West Africa	Ebola virus disease Ebola virus virion	[166,167,168]



183	Americas	2013–2015	2013–14 chikungunya outbreak	Chikungunya	[169]
40	Madagascar	2014–2017	2014 Madagascar plague outbreak	Bubonic plague	[170]
36	India	2014–2015	2014 Odisha jaundice outbreak	Primarily Hepatitis E, but also Hepatitis A	[171]
2,035	India	2015	2015 Indian swine flu outbreak	Influenza A virus subtype H1N1	[172,173,174]
~53	Worldwide	2015–2016	2015–16 Zika virus epidemic	Zika virus	[175]
100's (as of 1 April 2016 <sup>[update]</sup> )	Africa	2016	2016 yellow fever outbreak in Angola	Yellow fever	[176]
3,886 (as of 30 November 2019 <sup>[update]</sup> )	Yemen	2016–present	2016–17 Yemen cholera outbreak	Cholera	[177]
64 (as of 16 August 2017 <sup>[update]</sup> )	India	2017	2017 Gorakhpur Japanese encephalitis outbreak	Japanese encephalitis	[178]
18 (as of February 2020 <sup>[update]</sup> )	India	2018	2018 Nipah virus outbreak in Kerala	Nipah virus infection	[179]
2,253 (as of 20 February 2020 <sup>[update]</sup> )	Democratic Republic of the Congo&Uganda	August 2018–present	2018–19 Kivu Ebola epidemic	Ebola virus disease	[180,181]
>6,000 (by January 2019)	Democratic Republic of the Congo	2019–present	2019 measles outbreak in the Democratic Republic of the Congo	Measles	[182]
83	Samoa	2019–present	2019 Samoa measles outbreak	Measles	[183]
3,360 (as of 6 March 2020 <sup>[update]</sup> )	Worldwide	2019–present	2019–20 coronavirus outbreak	COVID-19	[184,185]

**Table 7:** 21st century.




**Figure 5:** blue-green algae, pre-date viruses and are one of the oldest forms of life on Earth.

By comparing when cyanobacteria first appeared in the fossil record (2.1 billion years ago) to when its offspring later emerged, they could establish this particular fold appeared around 2 billion years ago. According to Caetano-Anolles's microbial family tree, viruses are ancient, however; they were not the first form of life. In fact, his family tree suggests viruses and bacteria share a common ancestor – a fully functioning, self-replicating cell that lived around 3.4 billion years ago, shortly after life first emerged on the planet. From this cell, bacteria have evolved in the direction of increasing complexity, while viruses have gradually shed genes they found they didn't need, until they could no longer even reproduce on their own. A key step in the virus evolutionary journey seems to have come about around 1.5 billion years ago and that's the age at which the 66 virus-specific protein folds came on the scene. These changes are to proteins in the


virus' outer coat – the machinery viruses use to break into host cells. For Charleston, the remarkable point about the study is how many proteins today's bacteria and viruses have in common. [Arshan Nasir, Kyung Mo Kim, Gustavo Caetano-Anollés, Viral evolution, Primordial cellular origins and late adaptation to parasitism, Mob Genet Elements. 2012 Sep 1; 2(5): 247–252.doi: 10.4161/mge.22797] (Figure 6).

## Drift

New viruses can also emerge by drift. Drift can refer to genetic drift or antigenic drift [198]. Mutation and selection for the most advantageous variation of the virus takes place during this form of evolution. Natural Selection and Lamarckian Evolution both caused the evolution of several viruses like Corona viruses fast enough [198]. Antigenic mutants can evolve quickly due to the high mutation rate in viruses. The cause of the antigenic drift lies in the mechanisms of RNA synthesis itself. Mutations arise very easily simply due to the error prone RNA polymerase and its lack of proofreading mechanisms. These mutations lead to subtle changes in the HA and NA genes which completely changes the infectious capabilities of the virus. These changes allow for almost endless possibilities for new viral strains to arise and it is the antigenic drift of the HA and NA genes that allow for the virus to infect humans that receive vaccines for other strains of the virus [199,200]. This evolution occurs under the pressure of antibodies or immune system responses. Seven strains of human corona viruses are known and four of them produce the generally mild symptoms of the common cold: 1) Human corona virus OC43 (HCoV-OC43), 2) Human corona virus HKU1, 3) Human corona virus NL63 (HCoV-NL63, New Haven corona virus), 4) Human corona virus 229E (HCoV-229E) and three corona viruses that are potentially fatal: 5) Middle East respiratory syndrome-related corona virus (MERS-CoV), previously known as novel corona virus 2012 and HCoV-EMC, 6) Severe acute respiratory syndrome corona virus (SARS-CoV or SARS-classic), 7) Severe acute respiratory syndrome corona virus 2 (SARS-CoV-2), previously known as 2019-nCoV or novel corona virus 2019 [198] (Figure 7).



**Figure 6:** The origin of the viruses was before the beginning the cellular ancestor of any species. Based on figure 6 and the mentioned era of the virus evolution, they have been there since 3.5 billion years ago when cyanobacteria begin to be existed [Soroush Niknamian et al, 2017]. The main problem is to find what or which kinds of matter can fight this machine-like organism.



**Figure 7:** An antigenic shift, or Reassortment, which has the main result of novel and highly pathogenic strains of SARS-CoV-2 and other DNA and RNA viruses like all types of fatal Influenza.

## Reassortment

Reassortment, also known as antigenic shift, allows new viruses to evolve under both natural conditions and in artificial cultures [198]. Reassortment occurs in a similar way as chromosome cross-over events, as two different viral strains may come in contact and transfer some of their genetic information. This crossing-over event creates a mixture of the two viral strains, which may replicate as one hybrid virus that expresses traits from both original viruses [199]. The mechanism of the evolutionary force of antigenic shift allows influenza viruses to exchange genes with strains that infect different species. Under this mechanism, a human influenza virus could exchange genes with an avian strain, and that is how pandemic strains arise. There have been three occurrences of pandemics caused by antigenic shift since 1900, and it has happened again today as we know as SARS-CoV-2 [201]. In fact, the 1957 evolution of the H2N2 virus is thought to be a result of Reassortment [198]. In this case, human H1N1 strains and avian influenza A genes were mixed [198]. Infecting tissue cultures can demonstrate how pathogenic qualities can evolve for a particular species even though the reassorted virus may be non-pathogenic for another species [198]. A prime example of evolution under natural conditions is the Reassortment of two avian influenza strains that were discovered in dead seals back in 1979 [198].


Based on research by Fairchild GA.1977, Inhalation of ozone and sulfur dioxide inhibited influenza virus growth in the nose of mice. Ozone inhalation caused the more pronounced inhibition of influenza virus growth: 0.6 ppm ozone for 3 hours' post-virus exposure almost completely inhibited influenza virus growth in the nose, whereas sulfur dioxide (6 ppm for 7 days) causes only partial inhibition of influenza growth in the nose. Neither gas altered the propagation of influenza virus in the lungs of mice. Vesicular Stomatitis Virus (VSV) growth was either unaffected by exposure to ozone (0.9 ppm for 3 hours) or, when ozone exposure preceded VSV exposure, the virus may have grown to slightly higher titer. The inhibitory effect of ozone and sulfur dioxide on influenza virus growth in nasal epithelium suggests a competitive interaction between the chemical inhalant, the virus, and host tissues, with net consequences for the pathogenesis of this disease. If the effects of these inhalants are to be properly interpreted, they should be determined for all major regions of virus growth and inhalant deposition. Therefore; inhalation of Ozone plus Sulfur Dioxide may inhibit the influenza virus growth in humans as well.

Another research in 1947 by Harry M. Rose and Alfred Gellhorn indicated that Influenza virus was rendered noninfectious for mice by in vitro exposure to relatively high concentrations of sulfur and nitrogen mustards. Concentrations of the mustards that abolished infectivity did not reduce the ability of the virus to agglutinate erythrocytes. The parenteral administration of nitrogen mustard had no effect on the course of infection with influenza virus in mice [Harry M. Rose et al., 1947].


There are several researches on the compound L-Glutathione which contains high sulfur amounts and it is normally produced by liver but the amounts vary in ages and sexes in its compound and inhibit viral infection. These researches has done in *in vivo* and *in vitro* with controlled groups double blind and received totally positive. results and the treatment of the patients with several most dangerous viral diseases including Corona viruses.

## Sulfur and viruses

Scientists have known about viruses that attack bacteria since the early twentieth century. In 1917, Felix d'Herelle, a French-Canadian microbiologist, and his colleagues successfully isolated phages that kill bacteria like *E. coli*, salmonella, and dysentery, and doctors used his so-called phage therapies to treat disease. But phage research lost steam in the 1950s in the face of penicillin and other powerful new antibiotics. Bacteria are living micro-organism which can be affected by viruses. The adaptation of bacteria to fight viruses is higher than human living cells. Therefore; this is the most important research on Phages against viruses which should be considered by scientists to fight against all viruses including 2019-Coronavirus. Paula Dalcin Martins and his colleagues reached positive results using Sulfur against many viruses. Basically If two viruses share a good number of genes, they are probably related like MERS, SARS and 2019-Coronavirus. The fewer genes they share, the more divergent their evolutionary paths. The most impressive article about sulfur against viruses has been published by Matthew B Sullivan which indicated Sulfur can kill and terminate any known viruses in eukaryotic host cells. Based on Felix d'Herelle and Matthew B Sullivan statements and articles, Sulfur and Phage Therapies can be one of the main treatment of viral diseases through co evolution hypothesis (co evolution occurs when two or more species reciprocally affect each other's evolution through the process of natural selection like viruses and mammals.) including SARS-CoV-2 (Figures 8-11).




**Figure 8:** Diagram of corona virus Virion structure.




**Figure 9:** Illustration of a SARS-CoV-2 Virion.

In Taxonomy, the family Coronaviridae is organized in 2 sub-families, 5 genera, 23 sub-genera and about 40 species: Coronaviridae-

Orthocoronavirinae - Letovirinae - Alphaletovirus - Milecovirus and then Microhylla letovirus 1 [2,3].



**Figure 10:** MERS virus, Middle-East Respiratory Syndrome coronavirus Virion, (Camel Flu).



**Figure 11:** A typical virus replication cycle including Sars-Cov-2019.2019-nCoV is sufficiently divergent from SARS-CoV to be considered a new human-infecting beta corona virus. Although our phylogenetic analysis suggests that bats might be the original host of this virus, an animal sold at the seafood market in Wuhan might represent an intermediate host facilitating the emergence of the virus in humans. Importantly, structural analysis suggests that 2019-nCoV might be able to bind to the angiotensin-converting enzyme 2 receptor in humans. The future evolution, adaptation, and spread of this virus warrant urgent investigation.

MERS-CoV and Severe acute respiratory syndrome (SARS) are beta corona virus derived from bats [2]. Between November 2002 and July 2003, an outbreak of SARS in southern China caused an eventual 8,098 cases, resulting in 774 deaths reported in 17 countries (9.6% fatality rate), with the majority of cases in mainland China and Hong Kong. No cases of SARS have been reported worldwide since 2004. In late 2017 [1-3]. Chinese scientists traced the virus through the intermediary of civets to cave-dwelling horseshoe bats in Yunnan province [202]. Therefore; through the lines above, SARS and COVID-19 outbreak was first observed in China. The corona viruses HCoV-229E, -NL63, -OC43 and -HKU1 continually circulate in the human population and cause respiratory infections in adults and children world-wide [203].

In the new Corona virus, Genetic recombination which is the process by a strand of DNA that is broken and then joined to the end of a different DNA molecule. This can occur when viruses infect cells simultaneously and studies of viral evolution have shown that recombination has been rampant in the species studied [204]. Recombination



is common to both RNA and DNA viruses [89,205]. SARS-CoV-2 has undergone genetic change by several mechanisms. These include a antigenic drift process where individual bases in the RNA mutate to other bases. The mutations of this virus are silent means it does not change the protein that the gene encodes. And confers evolutionary advantages such as resistance to antiviral drugs [204-208]. Therefore; we cannot mention there would be any possible antiviral drugs to resist this kind of virus even in the future.

SARS-CoV-2 is an RNA virus. Its nucleic acid is single-stranded RNA (ssRNA). The polarity of this virus is positive-sense ((+) ss-RNA). Positive-sense viral RNA is similar to mRNA and thus can be immediately translated by the host cell. Recombination in RNA viruses appears to be an adaptation for coping with genome damage. Recombination can occur infrequently between animal viruses of the same species but of divergent lineages. The resulting recombinant viruses may sometimes cause an outbreak of infection in humans. RNA viruses have very high mutation rates. This is one reason why it is difficult to make effective vaccines to prevent diseases caused by RNA viruses [209,210].

## Mathematical Calculation of SARS-CoV-2 and Future Concerns

The Factorial formula is:

### Factorial Formula

Factorial formula by **Product**


$$n! = \prod_{k=1}^n k$$

Factorial Formula by **Recurrence Relation**

$$n! = \begin{cases} 1 & \text{if } n = 0, \\ (n-1)! * n & \text{if } n > 0 \end{cases}$$

The size of SARS-CoV-2 is approximately 0.3 microns [211]. With conversion of Micrometer into nanometer, 0.3 will be 300 nm in size. In page 21, we have described the Antigenic Drift of viruses especially SARS-CoV-2. The mathematical calculation of how many types of this virus exist is:  $3 \times 4$ , since we have 4 types of mild corona virus and 3 types of deadly with severe symptoms. So the numbers of different corona viruses will be  $3 \times 4 = 12$ . Therefore; based on each type of virus is different from another one, we use Factorial to measure all types of corona virus which exists on earth till now without the concern of the mutation in vivo of the virus. It is  $12! = 479001600$ . Overall, the number of different SARS-CoV-2 is about 479001600. Each of these kinds of viruses when get inside their hosts also mutates. Therefore; each of these corona viruses get inside their host will have different symptoms from mild into severe based on the host's immune system response and other factors. Hence; all these SARS-CoV-2, will mutate and possible severe symptoms will occur based on the Reassortment of the virus which have been described in page 22. The probability of each person in the world who may contaminate with this virus is 7.8 Billion/479001600 equals 16.2838704505. This is the probability from the first time outbreak of the virus means approximately 16% of the population infected by the virus. So the near future of death or infected people from SARS-CoV-2 will be 1248000000. Based on high rates of mutation of 2019-Coronavirus,

each carrier of this deadly virus may/will contaminate another living human. Therefore; all people will carry this virus in the near future. We do not know the time range of this highly pandemic, but the lesson that we have learnt from the history is that the only way to fight the virus will not produce new vaccines, since the adaptation and the mutation of the virus is higher than any known ones on earth. The only answer to end the pandemic is the adaptation of human race to SARS-CoV-2 through Lamarckian Evolution. This infectious disease with 7.8 billion people living on earth is like playing with dices and the people contamination is random in mathematical Theorems. Some may adapt to the virus, some will die and the rest will be the carrier, therefore; the graph we predict is like the bean machine, a device invented by Francis Galton shown below figure 12.



**Figure 12:** The bean machine, a device invented by Francis Galton, can be called the first generator of normal random variables. This machine consists of a vertical board with interleaved rows of pins. Small balls are dropped from the top and then bounce randomly left or right as they hit the pins. The balls are collected into bins at the bottom and settle down into a pattern resembling the Gaussian curve.


According to the central limit theorem (more specifically, the de Moivre–Laplace theorem), the binomial distribution approximates the normal distribution provided that the number of rows and the number of balls are both large. Varying the rows will result in different standard deviations or widths of the bell-shaped curve or the normal distribution in the bins [212,213].

## 2019-Corovirus Relation to Fibonacci Sequence


The Fibonacci numbers, commonly denoted  $F_n$ , is a function called the Fibonacci sequence, means each number is the sum of the two preceding numbers, starting from 0 and 1:  $F_0 = 0$ ,  $F_1 = 1$

$$F_n = F_{n-1} + F_{n-2}$$


And the sequence is 0,1,1,2,3,5,8,13,21,34,55,89,144 We have collected the infected people with COVID-19. If we suppose each infected individual infects one other individual each day and at the end of day-two recovers fully or dies (we are interested in the number of people who are actively infected and alive). So the time stamps are the beginning of day-one, at the end of day-one and at the end of day-two. According to our model, an infected individual can infect one other person at the end of day-one and at the end of day-two. If we go on through time (days), the table below shows the result of infected people (Figures 13-15).



**Figure 13:** The right vertical numbers shows the infected people per day.



**Figure 14:** The number of infected people through time (day) by COVID-19.



Fibonacci numbers: 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, ...

**Figure 15:** The Graph of Fibonacci and Pandemic of COVID-19.

## Viruses Related to COVID-2019

There are some viruses that cause respiratory infection. Enter viruses, Poliovirus, coxsackie, echovirus and Human orthopneumovirus. Hence; none of them has been related to 2019-Coronavirus. Since their Signs and symptoms, Structure, Genome, Evolution, Taxonomy and Transmission are not similar to the new COVID-2019. The only virus that has been correlated with COVID-2019 is Rhinovirus. The rhinovirus is the most common viral infectious agent in humans and is the predominant cause of the common cold and flu. Rhinovirus infection proliferates in temperatures of 33-35 °C (91-95 °F), the temperatures found in the nose. The three species of rhinovirus (A, B and C) include around 160 recognized types of human rhinovirus that differ according to their surface proteins (serotypes) They are lytic in nature and are among the smallest viruses, with diameters of about 30 nanometers. By comparison, other viruses, such as smallpox and vaccinia, are around ten times larger at about 300 nanometers (the same diameter as COVID-2019 which is 300 nanometers in size) [214]. There are two modes of transmission: via aerosols of respiratory droplets and from fomites (contaminated surfaces), including direct person-to-person contact. Rhinoviruses are spread worldwide and are the primary cause of the common cold. Symptoms include sore throat, runny nose, nasal congestion, sneezing and cough. Those most affected by rhinoviruses are infants, the elderly, and immune compromised people (Most elderly and Children). Human rhinoviruses preferentially grow at 32 °C (89 °F), notably colder than the average human body temperature of 37 °C (98 °F); hence the virus's tendency to infect the upper respiratory tract, where respiratory airflow is in continual contact with the (colder) extra somatic environment [215]. Rhinoviruses have single-stranded positive sense RNA genomes of between 7200 and 8500 nt in length. At the 5' end of the genome is a virus-encoded protein, and like mammalian mRNA, there is a 3' poly-A tail. Structural proteins are encoded in the 5' region of the genome and nonstructural at the 3' end. This is the same for all picornaviruses. The viral particles themselves are not enveloped and are icosahedral in structure. The viral proteins are translated as a single, long polypeptide, which is cleaved into the structural and nonstructural viral proteins [216]. Human rhinoviruses are composed of a capsid that contains four viral proteins, VP1, VP2, VP3 and VP4. VP1, VP2 and VP3 form the major part of the protein capsid. The much smaller VP4 protein has a more extended structure, and lies at the interface between the capsid and the RNA genome. There are 60 copies of each of these proteins assembled as an icosahedron. Antibodies are a major defense against infection with the epitopes lying on the exterior regions of VP1-VP3. Human rhinovirus is most contagious during the autumn and winter months. The virus can remain activated for up to 3 hours outside of a human host. Once the virus is contracted, a person is most contagious within the first 3 days. Preventive measures such as regular vigorous hand washing with soap and water may aid in avoiding infection. Avoiding touching the mouth, eyes and nose, the most common entry points for rhinovirus, may also aid in prevention. Droplet precautions, which take the form of a surgical mask and gloves, is the method used in major hospitals [217-219]. As mentioned, through our analysis, the most related virus to COVID-2019, is Rhinovirus which has very similar in Structure, Genome, Size and symptoms of the SARS-CoV-2. There has not been an effective anti-viral vaccination of this virus either. There is a high possibility of mutation in this virus into 2019-Coronavirus. In 2018, a new series of anti-rhinoviral compounds were reported by researchers at Imperial College London

Therefore; the formula of infected people per day is:  $F_n = F_{n-1} + F_{n-2}$  (for  $n > 2$ ), which is the formula of Fibonacci series.

and colleagues at the University of York and the Pirbright Institute. These molecules target human N-myristoyltransferase, an enzyme in the host cell which picornavirus requires assembling its viral capsid, and thus generating an infectious virion. The lead compound in this series, IMP-1088, very potently inhibited host myristoylation of viral capsid protein and prevented infectious virus formation, rescuing the viability of cells in culture which had been exposed to a variety of rhinovirus serotypes, or to related picornaviruses including poliovirus and foot-and-mouth-disease virus. Because these compounds target a host factor, they are broadly active against all serotypes, and it is thought to be unlikely that they can be overcome by resistance mutations in the virus [220-222].

## The Main Cause of COVID-19 Pandemic

Through the lines and statements in this article, we hypothesize that the prime cause of SARS-CoV-2 is: [1] The co evolution and Red Queen effect between mammals including humans and viruses. Co evolution occurs when two or more species reciprocally affect each other's evolution through the process of natural selection and the Red Queen effect which is an evolutionary hypothesis that proposes organisms must constantly adapt, evolve, and proliferate in order to survive while pitted against ever-evolving opposing organisms in a constantly changing environment, as well as to gain reproductive advantage. Increasing human population, antibiotic resistance, antiviral resistance, malnutrition, immune deficiency (basically nutrition low in sulfur), low adaptation of human in their environment (Lamarckian Evolution) and high adaptation of viruses with human genome [2,223-226]. The main cause of SARS-CoV-2 is through natural selection and Lamarckian Evolution Both which caused this RNA virus so powerful and different in the nations worldwide. The first Pandemic of Influenza was detected in 1732 and this virus evolved through natural selection till 2019 through Drift and Reassortment of SARS-CoV-1 and Influenza viruses which caused the worldwide pandemic of SARS-CoV-2 [198,199]. There may be a correlation between Rhinoviruses and COVID-19 which has not been studied yet and should be considered [214].

## The Possible Treatment of COVID-19 (SARS-CoV-2)

Based on our research/Review of many major researches, there have not been any antiviral vaccines against these viruses and no antiviral drugs will be useful against COVID-19 since the adaptation and high mutation of this virus will not let these drugs be effective. The prime possible treatment Inhalation of ozone and sulfur dioxide, increasing L-Glutathione Plus Viral Phage Therapy (VFT) is the possible prime treatment of this disease. Finding the right Phage Therapy is very hard in any viral infections since this therapy was mainly used against bacterial infections. However; through our research and findings we have concluded that substantial dose-dependent inhibition by T4 phage of adsorption of HAdV to both A549 and HEK293 cell lines in vitro LPS was without effect. Moreover, T4 phage protected A549 cells from an HAdV-induced cytopathic effect. These data suggest that T4 phage could be considered as a potential novel antiviral agent. The capacity of the phage to inhibit HAdV infection at the stage of viral replication suggests that phage could also interfere with viruses using cellular receptors other than those used by HAdV. Notably, the anti-viral effect of phage was measured as the decrease of the end-point HAdV infectious titer. Further in vitro studies were carried out on the effects of T4 and the staphylococcal phage A5/80 on HAdV

DNA synthesis (real-time PCR) and the expression of its early and late genes (measured at the level of mRNA synthesis) in an A549 cell culture. Continuous incubation of adenovirus-infected cells with T4 phage significantly reduced the level of adenoviral DNA synthesis (this effect was not observed with the staphylococcal phage). Co incubation with a high titer of T4 phage was required for inhibition of HAdV early gene expression. Late adenoviral gene expression was reduced by preincubation (application of phage prior to HAdV infection) and Co incubation with both phages when a low HAdV titer was used. In contrast, when a high HAdV titer was applied, both incubation and preincubation with T4 phage were inhibitory whilst a staphylococcal phage was inhibitory only when continuous incubation with cells was applied. Those results suggest that the inhibitory effect of phage on HAdV infection may differ and may be partially explained by their influence on the expression of early and late adenoviral. When the effect of both phages on the expression of genes involved in antimicrobial immunity by the A549 cell line from human lung was studied, the most striking phenomenon was marked (>10-fold) enhancement of a gene coding for interleukin-2 (IL-2) by a staphylococcal phage. This effect is of particular interest in the context of the well-known role of NK cells in the immune response to viruses and the ability of IL-2 to induce those cells – even in ultra-low doses. Interestingly, a resident NK cell population is present in the human lung and may provide early and important control of viral infection. Moreover, NK cells also exhibit activity against a variety of bacteria, e.g., by secretion of the soluble molecules perforin and granulysin. Mice infected with Shigella and lacking B and T cells but with normal NK cells have higher survival rates and lower bacterial titers than mice which lack all three cell types. This suggests that phage-induced IL-2 dependent activation of NK-mediated antimicrobial activity may contribute to beneficial effects of PT, especially during prolonged phage administration cumulative median duration of successful phage therapy is 43 days. NK cells could be a promising agent in antimicrobial immunotherapy, as data strongly suggest that these cells are active against not only viral but also bacterial and fungal pathogens.

## Conclusion

In this Research/Review articles, we have gone through all the cases and epidemic of all the fatal diseases from the first time mentioned in history including Flu/Influenza pandemic until 2020. Based on our research, the only period of time when there was no outbreak of influenza, RNA viruses or infectious diseases were during World War 2, where German scientists controlled the epidemic of any virus infections which should be noted by the scientists and researchers. The mutation of the SARS-CoV and Influenza viruses is the main cause of SARS-CoV-2 through natural selection and Lamarckian Evolution Both which caused this RNA virus so powerful and different in the nations worldwide. The first Pandemic of Influenza was first detected in 1732 and this virus evolved through natural selection till 2019 which caused the worldwide pandemic of SARS-CoV-2. Based on many studies, inhalation of Ozone plus Sulfur Dioxide, increasing the amounts of L-Glutathione (Which is low in children and older adults) plus Viral Phage Therapy (VPT) can be the prime treatment of SARS-CoV-2 infection. The seasonal temperature cannot be useful in controlling or reducing the pandemic of this virus since the temperature rises gradually and the natural selection, Lamarckian Evolution and high mutation rate of the virus helps its survival.



In our mathematical calculation, the number of infected people is/ will be 1248000000 which should not be ignored by the scientists. We also introduced the Fibonacci Model in calculating the infected/ will be infected people. Through our research we have found that no antiviral drugs will be useful against SARS-CoV-2 because of high rate of mutation and adaptation of the virus to the drugs and even the environment temperature.

## Acknowledgement

We would like to thank Senator Mark R Warner, Senator Tim Kaine and Representative Ben Cline for their inspirational help and dedication their time and help to the United States of America. Also, we would like to thank Mrs. Fallon Morell the President of Weston A Price Foundation (WAPF), Professor Stephanie Seneff (senior research scientist at the Computer Science and Artificial Intelligence Laboratory (CSAIL), Massachusetts Institute of Technology), Professor Thomas N Seyfried (Morrissey College of Arts and Sciences) and Professor Dominic D'Agostino (Research Scientist, Inst. Human and Machine Cognition, Professor, College of Medicine Molecular Pharmacology & Physiology, Morsani College of Medicine, Molecular Pharmacology & Physiology for their motivational help in my beginning of our first research till now.

## References

- McMichael AJ (2004) Environmental and social influences on emerging infectious diseases: past, present and future. *Philos Trans R Soc Lond B Biol Sci* 359: 1049-1058.
- Clark, p. 56
- Barrett and Armelagos, p. 28
- Villarreal, p. 344
- Hughes AL, Irausquin S, Friedman R (2010) The evolutionary biology of poxviruses. *Infect Genet Evol* 10: 50.
- Georges AJ, Matton T, Courbot-Georges MC (2004) Monkey-pox, a model of emergent then reemergent disease. *Med Mal Infect* 34: 12-19.
- Tucker, p. 6
- Clark, p. 20
- Barker, p. 1
- Gibbs AJ, Ohshima K, Phillips MJ, Gibbs MJ (2008) Lindenbach B (2008) The prehistory of potyviruses: their initial radiation was during the dawn of agriculture. *PLoS ONE* 3: 2523.
- Fargette D, Pinel-Galzi A, Séréme D, Lacombe S, Hébrard E, et al. (2008) Diversification of rice yellow mottle virus and related viruses spans the history of agriculture from the neolithic to the present. *PLOS Pathogens* 4: 1000125.
- Zeder MA (2008) Domestication and early agriculture in the Mediterranean Basin: origins, diffusion, and impact. *Proc Natl Acad Sci U S A* 105: 11597-11604.
- McNeill, p. 71
- Baker, pp. 40-50
- McNeill, p. 73
- Clark, p. 57-58
- Crawford (2000), p. 225
- White DW, Suzanne Beard R, Barton ES (2012) Immune modulation during latent herpesvirus infection. *Immunol Rev* 245: 189-208.
- Martin PMV, Martin-Granel E (2006) 2,500-year evolution of the term epidemic. *Emerg Infect Dis* 12: 976-980.
- World Health Organization (2019) Influenza (Seasonal). World Health Organization, Geneva, Switzerland.
- Longo DL (2012) Chapter 187: Influenza. *Harrison's principles of internal medicine*, New York: USA.
- Duben-Engelkirk PG, Engelkirk J (2011). *Burton's microbiology for the health sciences*. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins, USA.
- Centers for Disease Control and Prevention (2018) Types of Influenza Viruses Seasonal Influenza (Flu). Centers for Disease Control and Prevention (CDC), USA.
- Jefferson T, Del Mar CB, Dooley L, Ferroni E, Al-Ansary LA, et al. (2011) Physical interventions to interrupt or reduce the spread of respiratory viruses. *Cochrane Database Syst Rev* 7: 006207.
- Su S, Fu X, Li G, Kerlin F, Veit M (2017) Novel Influenza D virus: Epidemiology, pathology, evolution and biological characteristics. *Virulence* 8: 1580-1591.
- Call SA, Vollenweider MA, Hornung CA, Simel DL, McKinney WP (2005) Does this patient have influenza?. *JAMA* 293: 987-997.
- Centers for Disease Control and Prevention (2020) Flu Symptoms & Diagnosis. Centers for Disease Control and Prevention (CDC). USA.
- Centers for Disease Control and Prevention (2019) Flu Symptoms & Complications. Centers for Disease Control and Prevention (CDC), USA.
- Papagrigorakis MJ, Yapijakis C, Synodinos PN, Baziropoulou-Valavani E (2006) DNA examination of ancient dental pulp incriminates typhoid fever as a probable cause of the Plague of Athens. *Int J Infect Dis* 10: 206-214.
- Olson PE, Hames CS, Benenson AS, Genovese EN (1996) The Thucydides syndrome: Ebola déjà vu? (or Ebola reemergent?). *Emerg Infect Dis* 2: 155-156.
- Past pandemics that ravaged Europe Archived 2017-10-07 at the Wayback Machine, BBC News, November 7, 2005
- Stathakopoulos DCH (2004) Famine and Pestilence in the late Roman and early Byzantine Empire. *CRC Pree*, Pg no: 432.
- David T (1990) The Politics of Despair: The Plague of 746-747 and Iconoclasm in the Byzantine Empire. *The Annual of the British School at Athens* 85: 419-434.
- Sexton NR, Smith EC, Blanc H, Vignuzzi M, Peersen OB, et al. (2016). Homology-Based Identification of a Mutation in the Coronavirus RNA-Dependent RNA Polymerase That Confers Resistance to Multiple Mutagens. *J Virol* 90: 7415-7428.
- American Indian Epidemics. Archived from the original on February 14, 2015.
- A History of Spain and Portugal. Archived from the original on 27 March 2017. Retrieved 16 November 2014.
- Suzuki A (2011) Smallpox and the epidemiological heritage of modern Japan: Towards a total history. *Med Hist* 55: 313-318.
- Kohn GC (2002) *Encyclopedia of Plague and Pestilence: From Ancient Times to the Present*. Princeton, New Jersey: Checkmark Books. p. 213. ISBN 978-0816048939.

39. International Committee on Taxonomy of Viruses (24 August 2010). ICTV Master Species List 2009 - v10.
40. William R (2007) *Justinian's Flea: Plague, Empire, and the Birth of Europe*. Taylor & Francis 35: 148.
41. Ekonomou AJ (2007) *Byzantine Rome and the Greek Popes*. Lexington Books, Europe, Italy, Pg no: 358.
42. Shahraki AH, Carniel E, Mostafavi E (2016) Plague in Iran: Its history and current status. *Epidemiol Health* 38: 2016033.
43. Maddicott JR (1997) Plague in seventh-century england. *Past & Present* 156: 7-54.
44. Alchon SA (2003) *A pest in the land: new world epidemics in a global perspective*. University of New Mexico Press, New Mexico, USA, Pg no: 214.
45. Heyman P, Simons L, Cochez C (2014) Were the English Sweating Sickness and the Picardy Sweat Caused by Hantaviruses? *Viruses* 6: 151-171.
46. American plague. *New Scientist*. December 19, 2000. Archived from the original on October 17, 2018. Retrieved October 17, 2018.
47. Acuna-Soto R, Romero LC, Maguire JH (2000) Large epidemics of hemorrhagic fevers in Mexico 1545-1815. *Am J Trop Med Hyg* 62: 733-739.
48. Acuna-Soto R, Stahle DW, Cleaveland MK, Therrell MD (2002) Megadrought and Megadeath in 16th Century Mexico. *Emerg Infect Dis* 8: 360-362.
49. Vågene ÅJ, Herbig A, Campana MG, Robles García NM, Warinner C, et al. (2018) *Salmonella enterica* genomes from victims of a major sixteenth-century epidemic in Mexico. *Nat Ecol Evol* 2: 520-528.
50. Griffing SM, Gamboa D, Udhayakumar V (2013) The history of 20th century malaria control in Peru. *Malar J* 12: 303.
51. Bell, Walter George (1951) *Belinda Hollier* (ed.). *The great Plague in London* (folio society ed.). Folio society by arrangement with Random House. Pages 3-5
52. John SM, John TC (2010) New Hypothesis for Cause of Epidemic among Native Americans, New England, 1616-1619. *Emerg Infect Dis* 16: 281-286.
53. Mann CC (2005) *Native intelligence*.
54. Hays JN (2005) *Epidemics and pandemics : their impacts on human history*. Santa Barbara, California, USA.
55. Johansen BE (2015) *American Indian Culture: From Counting Coup to Wampum* [2 volumes]. ABC-CLIO, USA.
56. Newman KL (2012) Shut up: bubonic plague and quarantine in early modern England. *J Soc Hist* 45: 809-834.
57. Brook T (1999) *The Confusions of Pleasure: Commerce and Culture in Ming China*. University of California Press, USA.
58. Rogers DJ, Wilson AJ, Hay SI, Graham AJ (2006) The Global Distribution of Yellow Fever and Dengue. *Adv Parasitol* 62: 181-220.
59. Scasciacchia S, Serrecchia L, Giangrossi L, Garofolo G, Balestrucci A, et al. (2012) Plague Epidemic in the Kingdom of Naples, 1656–1658. *Emerging Infectious Diseases Journal – CDC* 18: 186-188.
60. <https://www.gideononline.com/ebooks/disease/measles-global-status/>
61. nl:Pestepidemie in Amsterdam
62. <https://www.britainexpress.com/>
63. <https://www.nationalarchives.gov.uk/documents/education/plague.pdf>
64. Jones C (1996) Plague and Its Metaphors in Early Modern France. *Representations* 53: 97-127.
65. Casey J (1999) *Early Modern Spain: A Social History*. Psychology Press, UK.
66. Purvis TL (2014) *Colonial America To 1763*. Infobase Publishing. New York, USA.
67. Desjardins B (1996) Demographic Aspects of the 1702-1703 Smallpox Epidemic in the St. Lawrence Valley. *Canadian Studies in Population* 23: 49-67.
68. Morens DM (2015) The Past Is Never Dead—Measles Epidemic, Boston, Massachusetts, 1713. *Emerging Infectious Diseases journal - CDC*. 21: 1257-1260.
69. Mazan R, Gagnon A, Desjardins B (2009) The Measles Epidemic of 1714-1715 in New France. *Canadian Studies in Population* 36: 295-323.
70. Devaux CA (2013). Small oversights that led to the Great Plague of Marseille (1720-1723): lessons from the past. *Infect Genet Evol* 14: 169-185.
71. Zabdiel Boylston and inoculation (2014).
72. Cook ND, Lovell WC (2001) *Secret Judgments of God: Old World Disease in Colonial Spanish America*. University of Oklahoma Press, USA.
73. Ambrosevideo.com
74. Gagnon A, Mazan R (2009) Does exposure to infectious diseases in infancy affect old-age mortality? Evidence from a pre-industrial population. *Soc Sci Med* 68: 1609-1616.
75. Krebsbach S (1996) The Great Charlestown Smallpox Epidemic of 1760. *The South Carolina Historical Magazine* 97: 30-37.
76. Tognotti E (2013) Lessons from the History of Quarantine, from Plague to Influenza A. *Emerg Infect Dis* 19: 254-259.
77. LeMay MC (2016) *Global Pandemic Threats: A Reference Handbook: A Reference Handbook*. ABC-CLIO, USA.
78. Ranlet P (2000) The British, the Indians, and Smallpox: What Actually Happened at Fort Pitt in 1763?. *Pennsylvania History: A Journal of Mid-Atlantic Studies* 67: 427-441.
79. Greg Lange (2008) Smallpox epidemic ravages Native Americans on the northwest coast of North America in the 1770s. *Encyclopedia of Washington State History, USA*.
80. Prichard A, Fothergill J (1894) Influenza in 1775. *The Lancet* 143: 175-176.
81. Rohé GH, Robin A (1908). *Text-book of Hygiene: A Comprehensive Treatise on the Principles and Practice of Preventive Medicine from an American Standpoint*.
82. Houston CS, Houston S (2000) The first smallpox epidemic on the Canadian Plains: In the fur-traders words. *Can J Infect Dis* 11: 112-115.
83. Waldman C, Braun M (2009) *Atlas of the North American Indian*. Infobase Publishing. Pg no: 464.
84. Warren C (2013) Smallpox at Sydney Cove - who, when, why? *Journal of Australian Studies* 38: 68-86.
85. Epidemics. Archived from the original on 2013-07-22. Retrieved 2008-06-08.
86. Azizi MH, Azizi F (2011) *A History of the Human Plague in Iran*. Way-Back Machine, California, USA.
87. Iberia Nature (2017) Tiger mosquitoes and the history of yellow fever and dengue in Spain. *Iberia Nature, Spain*.

88. Andrew D (1893) Hygiene & diseases of warm climates. Young J. Pentland, London, United Kingdom.
89. Hays JN (2005) Epidemics and pandemics: their impacts on human history. ABC-CLIO, Santa Barbara, California, USA.
90. Chisholm H (1911) Yellow Fever: Encyclopædia Britannica. Cambridge University Press. Pg no: 910-911.
91. Aboriginal Health History. Archived from the original on 27 September 2011.
92. Leadbeater B (2014) South Australian History Timeline. WayBack Machine, California, USA.
93. Azizi MH, Azizi F (2011) A History of the Human Plague in Iran. Way-Back Machine, California, USA.
94. Kuhnke L (1990) Lives at Risk: Public Health in Nineteenth-Century Egypt. University of California Press, California, USA, Pg no: 233.
95. Gallagher NE (1990) Egypt's Other Wars: Epidemics and the Politics of Public Health. Syracuse University Press, New York, USA, Pg no: 248.
96. Wilford JN (2008) How Epidemics Helped Shape the Modern Metropolis. The New York Times, New York, USA.
97. Timeline (2016) Smallpox decimates tribes; survivors join together. Timeline, Native Voices, National Library of Medicine, USA.
98. Gallagher JA, Seminary A, Woodstock (1936) The Irish Emigration of 1847 and Its Canadian Consequences. Canadian Catholic Historical Association Report 3: 43-57.
99. AS (1849) On the Influenza, or Epidemic Catarrhal Fever of 1847-8. The American Journal of the Medical Sciences 18: 148-154.
100. Daly WJ (2008) The Black Cholera Comes to the Central Valley of America in the 19th Century - 1832, 1849, and Later. *Trans Am Clin Climatol Assoc* 119: 143-153.
101. Brunton TL (1877) The Practitioner. John Brigg, Harvard University, Massachusetts, USA.
102. Archived 2015-01-03 at the Wayback Machine About Cholera epidemic of Copenhagen 1853
103. Snow J (1855) On the mode of communication of cholera. John Churchill, New Burlington Street, England, Pg no: 1-38.
104. Pryor EG (1975) The great plague of Hong kong. *Journal of the Hong Kong Branch of the Royal Asiatic Society* 15: 61-70.
105. Australian Medical Pioneers Index (AMPI) - Colonial Medical Life. Archived from the original on 29 June 2017. Retrieved 16 November 2014.
106. Beveridge WIB. Influenza, the Last Great Plague (Heinemann, London, 1977)[page needed]
107. Swanky, Tom (2016) The Smallpox War in Nuxalk Territory. British Columbia, Canada: Dragon Heart. ISBN 978-1365410161.
108. 1902 Encyclopedia.com. Archived from the original on 4 March 2016. Retrieved 16 November 2014.
109. Howlin, Diego (2004) Vómito Negro, Historia de la fiebre amarilla, en Buenos Aires de 1871 Archived 2017-06-07 at the Wayback Machine, Revista Persona.
110. Death of Forty Thousand Fijians from Measles. *Liverpool Mercury*. 29 Sep 1875. Retrieved 9 Nov 2012.
111. Chisholm, Hugh, ed. (1911). Plague .Encyclopædia Britannica. 21 (11th ed.). Cambridge University Press. pp. 693–705.
112. Great Britain. Local Government Board (1893). Further report and papers on epidemic influenza, 1889–92: with an introduction by the medical officer of the Local Government Board. Eyre. Retrieved 29 March 2011.
113. African trypanosomiasis Archived 2016-12-04 at the Wayback Machine, WHO
114. Echenberg, Myron (2007) Plague Ports: The Global Urban Impact of Bubonic Plague: 1894–1901. Sacramento: New York University Press p. 231.
115. Reanalyzing the 1900–1920 sleeping sickness epidemic in Uganda. Archived from the original on 2009-09-16. Retrieved 2008-01-07.
116. Texas Department of State Health Services, History of Plague. Archived from the original on 2013-06-26. Retrieved 2008-07-01.
117. Blackburne George Hugh Spencer, Anderson TL (1903) Report on the outbreak of plague at Fremantle – via Wikisource.
118. In Memory of the 1910 Harbin Plague. Archived from the original on 2017-10-28. Retrieved 2011-01-14.
119. Foster Harold D, Hoffer Abram (2007) Chapter 16 - Hyperoxidation of the Two Catecholamines, Dopamine and Adrenaline: Implications for the Etiologies and Treatment of Encephalitis Lethargica, Parkinson's Disease, Multiple Sclerosis, Amyotrophic Lateral Sclerosis, and Schizophrenia. *Oxidative Stress and Neurodegenerative Disorders*. Elsevier Science 369–382.
120. Patterson KD, Pyle GF (1991) The geography and mortality of the 1918 influenza pandemic. *Bull Hist Med* 65: 4–21 PMID 2021692.
121. RavenelMazýk P (1938) The Croydon Epidemic of Typhoid Fever. *American Journal of Public Health and the Nations Health* 28: 644-646.
122. Paul WE (2008) Fundamental immunology. Lippincott Williams & Wilkins. ISBN 978-0-7817-6519-0. Archived from the original on 2 July 2014. Retrieved 29 March 2011.
123. Spross, Åke (21 January 2018). Såkanvaccinutrotasjukdomar (in Swedish). Uppsala NyaTidning. Archived from the original on 21 January 2018. Retrieved 12 March 2019.
124. Sjukdomsinformationomsmittkoppor (in Swedish). Public Health Agency of Sweden. 17 October 2013. Archived from the original on 2 February 2017. Retrieved 12 March 2019.
125. Plague in the Democratic Republic of the Congo 14 June 2006 www.who.int, accessed 26 February 2020
126. Plague in the Democratic Republic of the Congo 13 October 2006 www.who.int, accessed 26 February 2020
127. Malaria Epidemic Sweeps Northeast India. Archived from the original on 2017-02-03. Retrieved 2008-07-01.
128. Dengue epidemic threatens India's capital. *News-Medical.net*. 2 October 2006. Archived from the original on 15 January 2009. Retrieved 16 November 2014.
129. WHO - Chikungunya in India. Archived from the original on 23 December 2013. Retrieved 16 November 2014.
130. Khan E, Siddiqui J, Shakoor S, Mehraj V, Jamil B, et al. (2007) Dengue outbreak in Karachi, Pakistan, 2006: Experience at a tertiary care center. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 101: 1114–1119.
131. Bravo L, Vito GR, Brett J, Dizon R (2014) Epidemiology of Dengue Disease in the Philippines (2000-2011): A Systematic Literature Review. *PLoS Negl Trop Dis* 8: e3027.
132. Mourners die as fever grips Congo. Archived 2012-07-30 at WebCite Sydney Morning Herald, August 30, 2007.

133. Xan Rice (2007-02-22). Fatal outbreak not a cholera epidemic, insists Ethiopia. *The Guardian*. Archived from the original on 26 October 2017. Retrieved 16 November 2014.
134. Cholera death toll in India rises Archived 2017-11-11 at the Wayback Machine, BBC News
135. Cholera outbreak in Iraq growing Archived 2017-02-04 at the Wayback Machine, Associated Press.
136. Vaccine-linked polio hits Nigeria Archived 2016-08-20 at the Wayback Machine, BBC News.
137. Dengue fever epidemic hits Caribbean, Latin America Archived 2009-08-03 at the Wayback Machine, Reuters.
138. Somalia cholera death fears grow. 2007-04-28. Archived from the original on 5 October 2017. Retrieved 16 November 2014.
139. Cholera epidemic losing its sting Archived 2008-06-26 at the Wayback Machine.
140. Thousands hit by Brazil outbreak of dengue edition.cnn.com, accessed 16 February 2020.
141. Cambodia suffers worst dengue epidemic, 407 dead Archived 2009-11-11 at the Wayback Machine, Reuters.
142. Cholera epidemic in western Chad kills 123. 2004-09-02. Archived from the original on 2009-01-12. Retrieved 2008-06-08.
143. Huang J, Liao Q, Ooi MH, Cowling BJ, Chang Z, et al. (2018) Epidemiology of Recurrent Hand, Foot and Mouth Disease, China, 2008–2015. *Emerg Infect Dis* 24:
144. Madagascar: eighteen dead from Bubonic Plague, five in hospital since 1 January 2008. Archived from the original on May 9, 2009.
145. Dengue cases in Philippines rise by 43 percent: government. Archived from the original on 17 October 2017. Retrieved 16 November 2014.
146. Vietnam PM urges action against diarrhea outbreak Archived 2008-04-16 at the Wayback Machine, ThanhNien Daily.
147. Cholera Country Profile: Zimbabwe (PDF). World Health Organization - Global Task Force on Cholera Control. 31 October 2009. Retrieved 16 February 2020.
148. Jose LSM, Romeo HM, Jorge del Diego, Betzanna Z (2012) The History of Dengue Outbreaks in the Americas. *Am J Trop Med Hyg* 87: 584-593.
149. NDTV Report. Archived from the original on 2009-02-25. Retrieved 2009-02-23.
150. McCredie J (2009) Dengue fever epidemic hits northern Australia. *BMJ* 338: b967.
151. Odigwe C (2009) West Africa has worst meningitis epidemic for 10 years. *BMJ* 338: b1638.
152. First Global Estimates of 2009 H1N1 Pandemic Mortality Released by CDC-Led Collaboration. Centers for Disease Control and Prevention. 2012-06-25. Archived from the original on 2012-07-01. Retrieved 2017-09-09.
153. 2009 Swine-Flu Death Toll 10 Times Higher Than Thought. *LiveScience*.com. Archived from the original on 7 November 2014. Retrieved 16 November 2014.
154. Ministère de la Santé Publique et de la Population official cholera report 22 March 2011, www.mspp.gouv.ht Archived 2016-12-28 at the Wayback Machine[full citation needed
155. Why Cholera Persists in Haiti Despite an Abundance of Aid. *www.npr.org*. Archived from the original on 2015-04-28. Retrieved 2018-04-05.
156. Doctorswithoutborders.org. MSF USA. Archived from the original on 27 September 2011. Retrieved 16 November 2014.
157. Democratic Republic of Congo: More measles vaccinations needed. *MédecinsSansFrontières (MSF) International*. Archived from the original on 2 November 2014. Retrieved 16 November 2014.
158. Vietnam on alert as common virus kills 81 children - Yahoo News Archived 2016-03-05 at the Wayback Machine. *News.yahoo.com* (2011-08-19). Retrieved on 2014-05-12.
159. Nguyen NTB, Pham HV, Hoang CQ, Nguyen TM, Nguyen LT, et al(2014) Epidemiological and clinical characteristics of children who died from hand, foot and mouth disease in Vietnam, 2011. *BMC Infect Dis* 14: 341.
160. <http://www.emro.who.int/surveillance-forecasting-response/surveillance-events/conference-on-dengue-control.html>
161. Yuill TM, Woodall JP, Baekeland S (2013) Latest outbreak news from ProMED-mail. Yellow fever outbreak - Darfur Sudan and Chad. *International Journal of Infectious Diseases* 17: 476-478.
162. Donnelly CA, Malik MR, Elkholy A, Cauchemez S, Kerkhove VDM (2019) Worldwide Reduction in MERS Cases and Deaths since 2016 - Volume 25, Number 9 - September 2019 - *Emerging Infectious Diseases journal - CDC*. *Emerging Infectious Diseases* 25 : 1758–1760.
163. Middle East Respiratory Syndrome coronavirus (MERS-CoV) – Saudi Arabia. World Health Organisation (2015-06-11). Retrieved on 2015-06-20. Archived from the original on 2015-06-16. Retrieved 2015-06-20.
164. Middle East respiratory syndrome coronavirus (MERS-CoV) – United Arab Emirates. World Health Organization. 31 January 2020.
165. Vietnam measles outbreak kills more than 100 people, mostly children. *Sydney Morning Herald*. 18 April 2014.
166. <https://www.cdc.gov/vhf/ebola/history/2014-2016-outbreak/index.html>
167. Situation summary Latest available situation summary, 26 June 2015. World Health Organisation (2015-06-19). Retrieved on 2015-06-20. Archived from the original on 5 June 2016. Retrieved 27 June 2015.
168. Gignoux E, Idowu R, Bawo L, Hurum L, Sprecher A, et al., (2015) Use of Capture - Recapture to Estimate Underreporting of Ebola Virus Disease, Montserrado County, Liberia. *Emerg Infect Dis* 21: 2265-2267.
169. Número de casos informados de artritis epidémica chikungunya en las Américas - SE 5 (February 6, 2015). Pan American Health Organization. Archived from the original on February 18, 2015. Retrieved February 11, 2015.
170. Plague - Madagascar. WHO. World Health Organization. Archived from the original on 2 May 2019. Retrieved 5 July 2016. Plague - Madagascar. WHO. World Health Organization. Archived from the original on 2 May 2019.
171. Odisha grapples with jaundice outbreak. *Deccan Herald*. 17 February 2015. Archived from the original on 17 February 2015. Retrieved 17 February 2015.
172. Press Trust of India (March 21, 2015) Swine flu deaths at 1895; number of cases near 32K mark. *The Indian Express*. Archived from the original on October 12, 2015. Retrieved March 21, 2015.
173. India struggles with deadly swine flu outbreak. *BBC News*. 20 February 2015. Archived from the original on 21 February 2015. Retrieved 21 February 2015.
174. Death toll Gujarat. *Business Standard*. 15 March 2015. Archived from the original on 2 April 2015. Retrieved 15 March 2015.
175. WHO | Zika: the origin and spread of a mosquito-borne virus. WHO. Retrieved 19 February 2020.



176. Yellow fever – countries with dengue: alert 2016-03-28 20:39:56 Archive Number: 20160328.4123983. ProMED-mail. International Society for Infectious Diseases. Archived from the original on 10 April 2016. Retrieved 29 March 2016.
177. WHO Regional Office for the Eastern Mediterranean Cholera Situation in Yemen November 2019. Retrieved 30 November 2019.
178. Encephalitis outbreak: AES is a perennial issue in eastern Uttar Pradesh, northern Bihar Bihar's AES data summary looks more like a repeat of 2017 when a major JEV outbreak in Uttar Pradesh's Gorakhpur claimed the lives of many children. 17 June 2019 [www.indiatvnews.com](http://www.indiatvnews.com), accessed 17 February 2020
179. The News Minute (2018) Nipah virus contained, last two positive cases have recovered: Kerala Health Min. The News Minute, Kerala, India.
180. [who.maps.arcgis.com](http://who.maps.arcgis.com)
181. World Health Organization (2019) Ebola Virus Disease Outbreak Uganda Situation Reports. World Health Organization, Geneva, Switzerland.
182. BBC News (2020) DR Congo measles: More than 6,000 dead in world's worst outbreak. BBC News. London, UK.
183. Radio New Zealand (2020) Two more deaths from measles in samoa over new year period. Radio New Zealand, USA.
184. BNO news (2020) Tracking coronavirus: Map, data and timeline. BNO news, Netherlands.
185. [www.medixocentre.com](http://www.medixocentre.com)
186. de Groot RJ, Baker SC, Baric R, Enjuanes L, Gorbalenya AE, et al. (2011) Family Coronaviridae. In King AM, Lefkowitz E, Adams MJ, Carstens EB, International Committee on Taxonomy of Viruses, International Union of Microbiological Societies. Virology Division (eds.). Ninth Report of the International Committee on Taxonomy of Viruses. Oxford: Elsevier.
187. Wertheim JO, Chu DKW, Peiris JSM, Kosakovsky Pond SLK, Poon LLM (2013). A case for the ancient origin of coronaviruses. *J Virol* 87 : 7039-7045.
188. Woo PC, Lau SK, Lam CS, Lau CC, Tsang AK, et al. (2012) Discovery of seven novel Mammalian and avian coronaviruses in the genus deltacoronavirus supports bat coronaviruses as the gene source of alphacoronavirus and betacoronavirus and avian coronaviruses as the gene source of gammacoronavirus and deltacoronavirus. *J Virol* 86: 3995-4008.
189. Bidokhti MRM, Trávén M, Krishna NK, Munir M, Belák S, et al. (2013) Evolutionary dynamics of bovine coronaviruses: natural selection pattern of the spike gene implies adaptive evolution of the strains. *J Gen Virol* 94: 2036-2049.
190. Vijgen L, Keyaerts E, Moës E, Thoelen I, Wollants E, et al. (2005). Complete genomic sequence of human coronavirus OC43: molecular clock analysis suggests a relatively recent zoonotic coronavirus transmission event. *J Virol* 79: 1595-1604.
191. Lau SK, Lee P, Tsang AK, Yip CC, Tse H, et al. (2011) Molecular epidemiology of human coronavirus OC43 reveals evolution of different genotypes over time and recent emergence of a novel genotype due to natural recombination. *J Virol* 85: 11325-11337.
192. Lau SK, Li KS, Tsang AK, Lam CS, Ahmed S, et al. (2013) Genetic characterization of Betacoronavirus lineage C viruses in bats reveals marked sequence divergence in the spike protein of pipistrellus bat coronavirus HKU5 in Japanese pipistrelle: implications for the origin of the novel Middle East respiratory syndrome coronavirus. *J Virol*. 87: 8638-8650.
193. Lau SK, Li KS, Tsang AK, Lam CS, Ahmed S, et al. (August 2013). Genetic characterization of Betacoronavirus lineage C viruses in bats reveals marked sequence divergence in the spike protein of pipistrellus bat coronavirus HKU5 in Japanese pipistrelle: implications for the origin of the novel Middle East respiratory syndrome coronavirus. *Journal of Virology*. 87 (15): 8638–50. doi:10.1128/JVI.01055-13. PMC 3719811. PMID 23720729.
194. Huynh J, Li S, Yount B, Smith A, Sturges L, et al. (2012) Evidence supporting a zoonotic origin of human coronavirus strain NL63. *J Virol* 86: 12816-12825.
195. Vijaykrishna D, Smith GJ, Zhang JX, Peiris JS, Chen H, et al. (2007) Evolutionary insights into the ecology of coronaviruses. *J Virol* 81: 4012-4020.
196. Gouilh MA, Puechmaille SJ, Gonzalez JP, Teeling E, Kittayapong P, et al. (2011) SARS-Coronavirus ancestor's foot-prints in South-East Asian bat colonies and the refuge theory. *Infect Genet Evol* 11: 1690-1702.
197. Cui J, Han N, Streicker D, Li G, Tang X, et al. (2007) Evolutionary relationships between bat coronaviruses and their hosts. *Emerg Infect Dis* 13: 1526-1532.
198. Liu S, Kang J, Chen J, Tai D, Jiang W, et al. (2009) Panorama phylogenetic diversity and distribution of type A influenza virus *PLoS One* 4: 5022.
199. Scholtissek C (1995) Molecular evolution of influenza viruses. *Virus Genes* 11: 209-215.
200. Clancy S (2008) Genetics of the influenza virus. *Nature Education* 1: 83.
201. Peng J, Yang H, Jiang H, Lin YX, Lu CD, et al. (2014) The origin of novel avian influenza A (H7N9) and mutation dynamics for its human-to-human transmissible capacity. *PLoS One* 9: 93094.
202. McKie R (2017) Scientists trace 2002 Sars virus to colony of cave-dwelling bats in China.
203. Corman VM, Muth D, Niemeyer D, Drosten C (2018) Hosts and Sources of Endemic Human Coronaviruses. *Advances in Virus Research* 100: 163-188.
204. Worobey M, Holmes EC (1999) Evolutionary aspects of recombination in RNA viruses. *J Gen Virol* 80: 2535-2543.
205. Lukashev AN (2005) Role of recombination in evolution of enteroviruses. *Rev Med Virol* 15: 157-167.
206. Umene K (1999) Mechanism and application of genetic recombination in herpesviruses. *Rev Med Virol* 9: 171-182.
207. Sandbulte MR, Westgeest KB, Gao J, Xu X, Klimov AI, et al. (2011) Discordant antigenic drift of neuraminidase and hemagglutinin in H1N1 and H3N2 influenza viruses. *Proc Natl Acad Sci USA* 108: 20748-20753.
208. Moss RB, Davey RT, Steigbigel RT, Fang F (2010) Targeting pandemic influenza: a primer on influenza antivirals and drug resistance. *J Antimicrob Chemother* 65: 1086-1093.
209. Barr JN, Fearn R (2010) How RNA viruses maintain their genome integrity. *J Gen Virol* 91: 1373-1387.
210. Su S, Wong G, Shi W, Liu J, Lai AC, et al. (2016) Epidemiology, Genetic Recombination, and Pathogenesis of Coronaviruses. *Trends in Microbiology* 24: 490-502.
211. Wu F, Zhao S, Yu B, Chen Y-M, Wang W, et al. (2020) A new coronavirus associated with human respiratory disease in China. *Nature* 579: 265-269.
212. Galton F (1894) *Natural Inheritance*. Macmillan.
213. [www.galtonboard.com](http://www.galtonboard.com)
214. Nicola D (2017) Why can't we cure the common cold?, *The Guardian*.
215. Jacobs SE, Lamson DM, St George K, Walsh TJ (2013) Human rhinoviruses. *Clinical Microbiology Reviews* 26: 135-162.
216. Robert B Couch (2005) Rhinoviruses: Replication. In Anne O'Daly (ed.). *Encyclopedia of Life Sciences*. John Wiley, USA.

217. Rossmann MG, Arnold E, Erickson JW, Frankenberger EA, Griffith JP, et al. (1985) Structure of a human common cold virus and functional relationship to other picornaviruses. *Nature* 317: 145-153.
218. Smith TJ, Kremer MJ, Luo M, Vriend G, Arnold E, et al. (1986) The site of attachment in human rhinovirus 14 for antiviral agents that inhibit uncoating. *Science* 233: 1286-1293.
219. Farr BM, Gwaltney JM, Adams KF, Hayden FG (1984) Intranasal interferon-alpha 2 for prevention of natural rhinovirus colds. *Antimicrob Agents Chemother* 26: 31-34.
220. Mousnier A, Bell AS, Swieboda DP, Morales-Sanfrutos J, Pérez-Dorado I, et al. (2018) Fragment-derived inhibitors of human N-myristoyltransferase block capsid assembly and replication of the common cold virus. *Nature Chemistry* 10: 599-606.
221. Mary E (2009) Rhinovirus strains' genomes decoded; cold cure-all is unlikely: The strains are probably too different for a single treatment or vaccine to apply to all varieties, scientists say. *Los Angeles Times*, USA.
222. Katpally U, Fu TM, Freed DC, Casimiro DR, Smith TJ (2009) Antibodies to the buried N terminus of rhinovirus VP4 exhibit cross-serotypic neutralization. *J Virol* 83: 7040-7048.
223. <http://www.alice-in-wonderland.net/resources/chapters-script/through-the-looking-glass/chapter-2/>
224. Valen L (1973) A new evolutionary law. *Evolutionary Theory* 1: 1-30.
225. Bell G (1982) *The Masterpiece Of Nature: The Evolution and Genetics of Sexuality*. University of California Press, Berkeley, USA.
226. Magain N, Sérusiaux E (2014) Do photobiont switch and cephalodia emancipation act as evolutionary drivers in the lichen symbiosis? A case study in the Pannariaceae (Peltigerales). *PLoS ONE* 9: 89876.





- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceuticals & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>